# **New Methods For Predictability Analysis**

Brian Farrell
Department of Earth and Planetary Sciences
Harvard University
Pierce Hall, 29 Oxford Street
Cambridge, MA 02138

phone: (617) 495-2998 fax: (617) 495-9837 email: farrell@deas.harvard.edu

Award #: N00014-99-1-0018

#### LONG-TERM GOAL

The long-term goal of this work is to improve atmospheric and oceanic deterministic forecast capability by maximizing the accuracy of the initial state estimate from which a forecast is made while minimizing observational and computational costs required to obtain the initial state.

# **OBJECTIVES**

The present objective of this work is to increase fundamental understanding of error dynamics and develop practical methods for computing error statistics. We intend to reduce the dimension of the error system so that error statistics can be practically obtained using forecast products. In addition from knowledge of the statistical structure of the time dependent error field we intend to determine the areas in space and time where observational resources can be most effectively applied.

### **APPROACH**

Small error dynamics is governed by linear equations but because these equations are non-normal and time dependent theoretical tools for analyzing error growth have only recently been obtained. Our approach is to apply these recent theoretical advances in non-normal time-dependent stability analysis to the forecast error problem. Methods of modern control theory, specifically balanced Hankel operator truncation methods, will be used to reduce the dimension of the error system.

### WORK COMPLETED

The fundamental theory of error growth in non normal time dependent forecast models has been developed.

The control theory approach of optimal balanced Hankel operator order reduction has been adapted and applied to model fluid error dynamics systems with encouraging results.

# **RESULTS**

Methods for obtaining the structure of the error field in a time dependent flow have been developed which facilitate prediction of error field statistics. We have shown that the dominant error growth arises from destabilization of a set of most non-normal vectors of the time mean operator, and that a very good basis for calculating the time dependent error is the first few evolved optimals of the time

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar DMB control number.	ion of information. Send comment arters Services, Directorate for Info	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	his collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 30 SEP 1999 2. RI		2. REPORT TYPE	3. DATES COVERED <b>00-00-1999 to 00-00-1999</b>		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
New Methods For Predictability Analysis				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Harvard University, Department of Earth and Planetary Sciences, 29 Oxford Street, Cambridge, MA, 02138				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAIL Approved for publ	ABILITY STATEMENT ic release; distributi	on unlimited			
13. SUPPLEMENTARY NO	OTES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	Same as Report (SAR)	4	

**Report Documentation Page** 

Form Approved OMB No. 0704-0188 mean propagator obtained over a period equal to the decorrelation time of the time varying flow (Fig. 1).




Fig. 1: Error structure for a model of the time dependent midlatitude barotropic jet as revealed by a histogram of projection of 500 observations of the error on the first three evolved optimals of the time mean flow. The evolved optimals considered are the left singular vectors of the propagator  $e^{\overline{A}T}$ , where  $\overline{A}$  is the linearized operator associated with time mean flow, and in this example T is the decorrelation time of the time dependent flow. The error projects primarily on these evolved optimals; projection on the remaining evolved optimals is negligible.

Methods for optimally reducing the dimension of the forecast error system have been applied to time independent error systems, as would be appropriate for forecast over a short enough interval that time dependence of the flow can be ignored (~24 hr in the atmosphere). This optimal reduction of order is obtained by truncating the error dynamics using a balanced representation in the stochastic optimals and the empirical orthogonal functions of the system (Moore, 1981; Glover, 1984; Farrell and Ioannou, 1996).

Methods for quantifying the accuracy of the reduced low order error system have been developed. These methods have been applied to a number of model time independent forecast error systems in which a very satisfactory approximation to the error dynamics was obtained with dimension reduction by a factor of 10 (Fig. 2).




Fig. 2: Maximum error growth as a function of time for a model of the midlatitude barotropic mean jet. The original system (solid) has 100 degrees of freedom and various truncations of order 6 are shown. The best results are obtained for the balanced truncation (circles). Also shown are the optimal growth for an order 6 approximate system in which the top 6 stochastic optimals (SO's) have been retained (crosses); the optimal growth for an order 6 approximate system in which the top 6 empirical orthogonal functions (EOF's) have been retained (stars); and the optimal growth for an order 6 approximate system in which the top 6 least damped modes have been retained (dash-dot). The graph demonstrates that even with 6 degrees of freedom accurate error evolution can be obtained using balanced truncation.

It was determined that truncating the dynamical system in a balanced realization of the optimals and evolved optimals for a single time provides nearly optimal reduced order error system so long as the optimals are obtained for growth over an interval corresponding to the global optimal time. This is a significant result because it suggests that implementing the order reduction algorithm in an operational forecast mode will not be computationally difficult.

#### IMPACT/APPLICATION

The methods developed in this work can be directly extended to operational forecast models. Efficient calculation of error statistics will allow implementation of advanced methods of initial state estimation such as the Kalman filter.

#### **TRANSITIONS**

None

### RELATED PROJECTS

None

# **REFERENCES**

Farrell, B. F. and Ioannou P. J. 1996: Generalized Stability. Part I: Autonomous Operators. Journal of the Atmospheric Sciences, 53, 2025-2041.

Glover, K. 1984: An optimal Hankel-norm approximation of linear multivariable systems and their  $L^{\infty}$  error bounds. International Journal of Control, 39, 1115-1193.

Moore, B. C., 1981: Principal component analysis in linear systems: controllability, observability, and model reduction. IEEE Trans. on Automatic Control, AC-26, 17-31.

# **PUBLICATIONS**

Farrell, B. F. and Ioannou, P. J. 1999: Perturbation growth and structure in time dependent flows. Journal of the Atmospheric Sciences. In Press.

Farrell, B. F. and Ioannou, P. J. 1999: Approximating an Autonomous Stable Linear Dynamical System by a System of Lower Order. Journal of the Atmospheric Sciences. Submitted.